arXiv:1605.00812v1 [math.PR] 3 May 2016

The Cameron-Martin Theorem for (p-)Slepian processes

Wolfgang Bischoff* **and Andreas Gegg**

Faculty of Mathematics and Geography,
Catholic University of Eichstätt-Ingolstadt,

Abstract

We show a Cameron-Martin theorem for Slepian processes $W_t := \frac{1}{\sqrt{p}}(B_t - B_{t-p}), t \in [p,1]$, where $p \geq \frac{1}{2}$ and B_s is Brownian motion. More exactly, we determine the class of functions F for which a density of $F(t) + W_t$ with respect to W_t exists. Moreover, we prove an explicit formula for this density. p-Slepian processes are closely related to Slepian processes. p-Slepian processes play a prominent role among others in scan statistics and in testing for parameter constancy when data are taken from a moving window.

Keywords: Cameron-Martin theorem, (p-)Slepian process, Radon-Nikodym derivative
subject classification: 60G15, 60H99

1 Introduction

A Cameron-Martin theorem for a stochastic process is one of the most useful tools to solve problems the process is involved in. Let a stochastic process $X_{[a,b]} = (X_t)_{t\in[a,b]}$ with paths in $C[a,b]$, the set of real-valued and continuous functions on $[a,b] \subseteq \mathbb{R}$, and a deterministic function $F \in C[a,b]$ be given. Let $\mathbb{P}^{X_{[a,b]}}$ and $\mathbb{P}^{(X+F)_{[a,b]}}$ denote the distribution of $X_{[a,b]}$ and of $(X_t + F(t))_{t\in[a,b]}$ on $C[a,b]$, respectively. Then a Cameron-Martin theorem gives conditions on F under which a density $\frac{d(\mathbb{P}^{X_t+F(t)})}{d\mathbb{P}^{X_t}}$ exists and, additionally, it gives an explicit formula for this density.

The first and best known result of this type is by [6]. They proved it for the standard Brownian motion $B_{[0,1]} := (B_t)_{t\in[0,1]}$ with continuous paths. This Cameron-Martin theorem can be used, for example, to calculate optimal tests, see [4], and to estimate boundary crossing probabilities, see [5]. Therefore, such a result of Cameron-Martin type is also of great interest for other stochastic processes. The following theorem, see Lifshits [12, Theorem 5.1], can be used as basis to get results of Cameron-Martin type for centered Gaussian processes. To this end, let $L^2(C[a,b],P)$ be the set of all real-valued and square-integrable

*corresponding author: Wolfgang Bischoff, Faculty of Mathematics and Geography, D-85071 Eichstätt, Germany; E-Mail: wolfgang.bischoff@ku-eichstaett.de

functions on $C[a,b]$ with respect to a measure P defined on the Borel-σ-Algebra of $C[a,b]$.

Theorem 1.1 (An abstract Cameron-Martin theorem)
Let $X_{[a,b]} := (X_t)_{t\in[a,b]}$ be a centered Gaussian process with paths in $C[a,b]$, let $\mathcal{H} = \mathcal{H}_{X_{[a,b]}} \subseteq C[a,b]$ be the kernel of $\mathbb{P}^{X_{[a,b]}}$ and $\|\cdot\|_{\mathcal{H}}$ its inherent norm. Then,

$$\mathbb{P}^{(X+h)_{[a,b]}} \text{ is absolutely continuous with respect to } \mathbb{P}^{X_{[a,b]}} \iff h \in \mathcal{H} .$$

If $h \in \mathcal{H}$, then

$$\frac{d\mathbb{P}^{(X+h)_{[a,b]}}}{d\mathbb{P}^{X_{[a,b]}}}(g) = \exp\left(-\frac{1}{2}\|h\|_{\mathcal{H}}^2 + z(g)\right) \quad \text{for } \mathbb{P}^{X_{[a,b]}}\text{-almost all } g \in C([a,b]),$$

where $z = z_{X_{[a,b]}} \in L^2(C[a,b], \mathbb{P}^{X_{[a,b]}})$ is a linear functional fulfilling the equation

$$\mathbf{E}\left(X_t \cdot z_{X_{[a,b]}}\right) = h(t) , \quad t \in [a,b].$$

Hence, the kernel $\mathcal{H} = \mathcal{H}_{X_{[a,b]}}$ of the stochastic process $X_{[a,b]}$ together with its inherent norm $\|\cdot\|_{\mathcal{H}}$ and the functional $z_{X_{[a,b]}}$ must be determined to obtain an applicable result.

In the following, we use the notation stated in the next definition.

Definition 1.2
Let $X_{[a,b]} := (X_t)_{t\in[a,b]}$ be a centered Gaussian process with paths in $C[a,b]$ and let $\mathcal{H} = \mathcal{H}_{X_{[a,b]}} \subseteq C[a,b]$ be the kernel of $\mathbb{P}^{X_{[a,b]}}$. Then, we say that a function

$$z = z_{X_{[a,b]}} : \mathcal{H}_{X_{[a,b]}} \times C[a,b] \to \mathbb{R}$$

fulfills condition A for $X_{[a,b]}$, if and only if for all $h \in \mathcal{H}_{X_{[a,b]}}$ the function $z(h,\cdot) \in L^2(C[a,b], \mathbb{P}^{X_{[a,b]}})$ is a linear functional fulfilling

$$\mathbf{E}\left(X_t \cdot z(h, X_{[a,b]})\right) = h(t) , \quad t \in [a,b]. \tag{1}$$

For the Brownian motion, we have the following result by [6]. To this end, let λ denote the Lebesgue measure on $\mathbb{R}$ and let $L^2[a,b] := L^2([a,b],\lambda)$ be the square-integrable functions on $[a,b]$ with respect to λ.

Theorem 1.3
The kernel of $B_{[0,1]}$ is given by

$$\mathcal{H}_{B_{[0,1]}} := \{s_f \in C[0,1] \mid f \in L^2[0,1]\},$$

where

$$s_f(t) := \int_{[0,t]} f \, d\lambda \, , t \in [0,1], \text{ for } \lambda\text{-integrable } f : [0,1] \to \mathbb{R}.$$

It is furnished with the norm

$$\|s_f\|_{\mathcal{H}_{B_{[0,1]}}} := \|f\|_{L^2[0,1]}, \quad f \in L^2[0,1].$$

A function $z := z_{B_{[0,1]}}$ fulfilling condition A for $B_{[0,1]}$ is given by the Wiener integral, see, for instance, [14],

$$z : \mathcal{H}_{B_{[0,1]}} \times C[0,1] \to \mathbb{R}, \quad (s_f, b) \to \int_0^1 f \, db.$$

Slepian and p-Slepian processes are defined and their close relations are discussed in the next section. In Sect. 3, results of Cameron-Martin type are established for p-Slepian processes. Several proofs are postponed to an appendix.

2 Slepian and p-Slepian processes

The Slepian process X is the centered stationary Gaussian process with covariance function

$$C_X(s', s'+u) = (1-u)^+ \quad , \; 0 \le s' \le s'+u,$$

where $t^+ = \max(0,t), t \in \mathbb{R}$. This process was introduced and studied in [18] and later in [16,17]. Afterward, it was handled in numerous theoretical and applied probabilistic models; see, e.g., [1-3,9-11,13,15].

Let us consider the Slepian process on time intervals $[1,b]$, where $1 < b$ is any fixed constant. Let ∇_p , $p > 0$, be the backward difference operator with lag p, i.e., $\nabla_p F(t) = F(t) - F(t-p), t \in \mathbb{R}$, for functions $F : \mathbb{R} \to \mathbb{R}$. Note that the Slepian process coincides in distribution with

$$(\nabla_1 B)_{[1,b]} := (\nabla_1 B_t)_{t\in[1,b]} = (B_t - B_{t-1})_{t\in[1,b]} \ ,$$

where B_t is the standard Brownian motion with continuous paths in $[0,b]$. In Cressie [8, pp. 834], a slightly different expression of Slepian processes appeared in connection with scan statistics. They were considered the processes

$$\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]} := \frac{1}{\sqrt{p}}(\nabla_p B_t)_{t\in[p,1]} = \frac{1}{\sqrt{p}}\,(B_t - B_{t-p})_{t\in[p,1]} \tag{2}$$

for any fixed constant $p \in (0,1)$. We call this process p-Slepian process. Another application of p-Slepian processes is given by [7]. They consider moving sums of recursive residuals which are taken from windows of length p. By letting the number of residuals to ∞, they get the p-Slepian process $\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]}$.

By a suitable scaling in time the Slepian process $(\nabla_1 B)_{[1,b]}, b \in (1,\infty)$, can be transferred to the p-Slepian process $\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]}, p \in (0,1)$, by putting $p = \frac{1}{b}$. More exactly it holds $(\nabla_1 B_u)_{u\in[1,b]} = \sqrt{b}(\nabla_{\frac{1}{b}} B_{\frac{u}{b}})_{u\in[1,b]}$ in distribution. The covariance function of a p-Slepian process is given by

$$C_{\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]}}(s, s+t) = \left(1 - \frac{t}{p}\right)^+ \quad , p \le s \le s+t \le 1.$$

3 Cameron-Martin Theorem for p-Slepian processes

The function $\frac{1}{\sqrt{p}}\nabla_p, 0 < p \leq 1$, is linear. Hence, by [12] proposition 4.1, the kernel $\mathcal{H}_{p-\mathrm{Sl}}$ of the p-Slepian process $\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]}$ is given by $\mathcal{H}_{p-Sl} = \frac{1}{\sqrt{p}}\nabla_p(\mathcal{H}_{B_{[0,1]}}) = \{\frac{1}{\sqrt{p}}\nabla_p h | h \in \mathcal{H}_{B_{[0,1]}}\}$. By some calculations we get

$$\begin{aligned}\mathcal{H}_{p-\mathrm{Sl}} &= \left\{\frac{1}{\sqrt{p}}\nabla_p s_f : [p,1] \to \mathbb{R} \mid f \in L^2[0,1]\right\} \\ &= \left\{c + s_g : [p,1] \to \mathbb{R} \mid c \in \mathbb{R}\ ,\ g \in L^2[p,1]\right\}.\end{aligned}$$

More exactly, we have for $f \in L^2[0,1]$

$$\frac{1}{\sqrt{p}}\nabla_p s_f(t) = \frac{1}{\sqrt{p}}s_f(p) + s_{\frac{1}{\sqrt{p}}\nabla_p f}(t), t \in [p,1]. \tag{3}$$

It is much more complicated to determine the inherent norm of $\mathcal{H}_{p-\mathrm{Sl}}$ since ∇_p is not injective. For the following, the information of $\nabla_p s_f$ is important. For $f, g \in L^2[0,1]$, it holds true

$$\forall t \in [p,1] : \nabla_p s_f(t) = s_f(p) + \int_p^t f(s) - f(s-p)ds = \nabla_p s_g(t) \Longleftrightarrow$$

$$f(t) - f(t-p) = g(t) - g(t-p)\ \lambda - a.s. \text{ for all } t \in [p,1],\quad s_f(p) = s_g(p).$$

We prove the following result in the appendix.

Lemma 3.1 *Let $1/2 \leq p \leq 1$. Then the kernel $\mathcal{H}_{p-Sl}$ is furnished with its inherent norm*

$$\begin{aligned}&\left\|\frac{1}{\sqrt{p}}\nabla_p s_f\right\|^2_{p-Sl} = \frac{1}{p}\inf_{g \in L^2[0,1]:\nabla_p s_f = \nabla_p s_g}\|g\|^2_{L^2[0,1]} \\ &= \frac{1}{2p(3p-1)}(2s_f(p) + \delta(1-p))^2 + \frac{1}{2p}\|(f(t) - f(t-p))_{t\in[p,1]}\|^2_{L^2[p,1]},\end{aligned} \tag{4}$$

where $\delta = \frac{1}{1-p}(s_f(1) - s_f(p) - s_f(1-p))$. This minimum is attained at the function

$$\begin{aligned}f^*(t) = &\left(\frac{1}{3p-1}(s_f(p) + \frac{1-p}{2}\delta) + \frac{1}{2}(-f(t+p) + f(t))\right)\mathbf{1}_{[0,1-p]}(t) \\ &+ \frac{1}{2p-1}(s_f(p) - \frac{1-p}{3p-1}(s_f(p) - (2p-1)\delta))\mathbf{1}_{(1-p,p)}(t) \\ &+ \left(\frac{1}{3p-1}(s_f(p) + \frac{1-p}{2}\delta) + \frac{1}{2}(f(t+p) - f(t))\right)\mathbf{1}_{[p,1]}(t), t \in [0,1].\end{aligned}$$

Finally, we need the function $z := z_{\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]}}$ fulfilling condition A for the p-Slepian process to be in the position to state a result of Cameron-Martin type.

Lemma 3.2 *Let $1/2 \leq p \leq 1$. The function $z := z_{\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]}}$ defined by*

$$z : \mathcal{H}_{\frac{1}{\sqrt{p}}(\nabla_p B)_{[0,1]}} \times C[p,1] \to \mathbb{R}, \quad \left(\frac{1}{\sqrt{p}} s_f(p) + s_{\frac{1}{\sqrt{p}}\nabla_p f}, \frac{1}{\sqrt{p}} \nabla_p b \right) \mapsto \tag{5}$$

$$\frac{3s_f(p) + s_{\nabla_p f}(1)}{2(3p-1)} \left(\nabla_p b(p) + \nabla_p b(1) \right) + \frac{1}{2} \int_p^1 \nabla_p f(s) \, d(\nabla_p b(s)) \tag{6}$$

fulfills condition A for $\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]}$. Note that the integral

$$\int_p^1 \nabla_p f(s) \, d(\nabla_p B(s)) = \int_p^1 \nabla_p f(s) \, dB(s) - \int_p^1 \nabla_p f(s) \, dB(s-p)$$

has to be understood as Wiener Integral.

The proof is given in the appendix. By the above lemmas, we obtain the following result.

Theorem 3.3 (Cameron-Martin theorem for p-Slepian processes) *Let $1/2 \leq p \leq 1$. It holds true*

$$\mathbb{P}^{(\frac{1}{\sqrt{p}}\nabla_p B + h)_{[p,1]}} \text{ is absolutely continuous with respect to } \mathbb{P}^{(\frac{1}{\sqrt{p}}\nabla_p B)_{[p,1]}}$$
$$\iff h \in \mathcal{H}_{p-Sl} \, .$$

For $h = \frac{1}{\sqrt{p}} s_f(p) + s_{\frac{1}{\sqrt{p}}\nabla_p f}(\cdot) \in \mathcal{H}_{p-Sl}$ it holds true for $\mathbb{P}^{(\frac{1}{\sqrt{p}}\nabla_p B)_{[p,1]}}$-almost all $\frac{1}{\sqrt{p}}\nabla_p b \in C[p,1]$:

$$\begin{aligned}
&\frac{d\mathbb{P}^{(\frac{1}{\sqrt{p}}\nabla_p B + h)_{[p,1]}}}{d\mathbb{P}^{(\frac{1}{\sqrt{p}}\nabla_p B)_{[p,1]}}} \left(\frac{1}{\sqrt{p}} \nabla_p b \right) \\
&= \exp\left(-\frac{1}{4p} \left(\frac{(2s_f(p) + \delta(1-p))^2}{3p-1} + \| (f(\cdot) - f(\cdot - p)) \|^2_{L^2[p,1]} \right) \right. \\
&\quad \left. + \frac{3s_f(p) + s_{\nabla_p f}(1)}{2(3p-1)} (\nabla_p b(p) + \nabla_p b(1)) + \frac{1}{2} \int_p^1 \nabla_p f(s) \, d(\nabla_p b(s)) \right),
\end{aligned}$$

where $\delta = \frac{1}{1-p}(s_f(1) - s_f(p) - s_f(1-p))$.

4 Appendix

Proof of Lemma 3.1 Let $\frac{1}{\sqrt{p}} \nabla_p s_f \in \mathcal{H}_{p-\mathrm{Sl}}, f \in L^2[0,1]$, be arbitrary and let $1/2 \leq p \leq 1$. There are several places where the case $p = 1/2$ must be considered separately. We do not do this since the specifications for $p = 1/2$ are simpler than the following considerations for $1/2 < p \leq 1$. By proposition 4.1 of [12], we get

$$\begin{aligned}
\| \frac{1}{\sqrt{p}} \nabla_p s_f \|_{p-Sl} &= \frac{1}{\sqrt{p}} \inf_{g \in L^2[0,1] : \nabla_p s_f = \nabla_p s_g} \| s_g \|_{\mathcal{H}_{B_{[0,1]}}} \\
&= \frac{1}{\sqrt{p}} \inf_{g \in L^2[0,1] : \nabla_p s_f = \nabla_p s_g} \| g \|_{L^2[0,1]},
\end{aligned}$$

where the second equality follows by Theorem 1.3. Each function $f \in L^2[0,1]$ can be written $\lambda - a.s.$ in the form

$$f(t) = \alpha \mathbf{1}_{[0,1-p]}(t) + a(t) + \beta \mathbf{1}_{(1-p,p)}(t) + b(t) + \gamma \mathbf{1}_{[p,1]}(t) + c(t), t \in [0,1], \quad (7)$$

where

$$\begin{aligned}
\alpha &= \frac{1}{1-p} s_f(1-p), a(t) = \mathbf{1}_{[0,1-p]}(t)(f(t) - \alpha), \\
\beta &= \frac{1}{2p-1}(s_f(p) - s_f(1-p)), b(t) = \mathbf{1}_{(1-p,p)}(t)(f(t) - \beta), \\
\gamma &= \frac{1}{1-p}(s_f(1) - s_f(p)), c(t) = \mathbf{1}_{[p,1]}(t)(f(t) - \gamma).
\end{aligned}$$

Note that the six summands in (7) build orthogonal functions in $L^2[0,1]$. In the following, we use the consequence of this representation at several places without citing it explicitly. Furthermore, we have

$$s_f(p) = \alpha(1-p) + \beta(2p-1), f(t) - f(t-p) = \gamma - \alpha + c(t) - a(t), t \in [p,1].$$

Next, we consider $g \in L^2[0,1]$ of the specific form

$$g(t) = \alpha \mathbf{1}_{[0,1-p]}(t) + \beta \mathbf{1}_{(1-p,p)}(t) + (\alpha + \delta)\mathbf{1}_{[p,1]}(t), t \in [0,1], \alpha, \beta, \delta \in \mathbb{R}.$$

Hence,

$$s_g(p) = \alpha(1-p) + \beta(2p-1) \Leftrightarrow \beta = \frac{1}{2p-1}(s_g(p) - \alpha(1-p))$$

and

$$g(t) - g(t-p) = \delta, t \in [p,1].$$

Thus, $\nabla_p(g)(t) = s_g(p) + \delta t, t \in [p,1]$. Let $g_0(t) = \alpha_0 \mathbf{1}_{[0,1-p]}(t) + \beta_0 \mathbf{1}_{(1-p,p)}(t) + (\alpha_0 + \delta_0)\mathbf{1}_{[p,1]}(t) \in L^2[0,1]$ be fixed and let $g \in L^2[0,1]$ with $\nabla_p(g)(t) = \nabla_p(g_0)(t) = s_{g_0}(p) + \delta_0 t, t \in [p,1]$. Then,

$$g(t) = \alpha \mathbf{1}_{[0,1-p]}(t) + \frac{1}{2p-1}(s_{g_0}(p) - \alpha(1-p))\mathbf{1}_{(1-p,p)}(t) + (\alpha + \delta_0)\mathbf{1}_{[p,1]}(t)$$

and the square of its norm is given by

$$\|g\|^2 = \frac{1-p}{2p-1}(\alpha^2(3p-1) - 2\alpha(s_{g_0}(p) - (2p-1)\delta_0)) + \frac{s_{g_0}(p)^2}{2p-1} + \delta_0^2(1-p).$$

This norm is minimal if and only if

$$\alpha = \frac{1}{3p-1}(s_{g_0}(p) - (2p-1)\delta_0).$$

Hence, we obtain after some calculation

$$\|\nabla_p s_{g_0}\|^2_{p-Sl} = \frac{1}{3p-1}(2s_{g_0}(p)^2 + 2(1-p)s_{g_0}(p)\delta_0 + \delta_0^2(1-p)p).$$

Next, we consider a general function

$$f = \alpha \mathbf{1}_{[0,1-p]}(t) + a(t) + \beta \mathbf{1}_{(1-p,p)}(t) + b(t) + (\alpha + \delta)\mathbf{1}_{[p,1]}(t) + c(t) \in L^2[0,1].$$

By the above considerations we obtain

$$\begin{aligned}\|\nabla_p s_f\|_{p-Sl}^2 =& \frac{1}{3p-1}(2s_f(p)^2 + 2(1-p)s_f(p)\delta + \delta^2(1-p)p) \\ &+ \min(\|a(t)\|^2 + \|b(t)\|^2 + \|c(t)\|^2)\end{aligned}$$

where the minimum is taken over all $a : [0, 1-p] \to \mathbb{R}, b : (1-p, p) \to \mathbb{R}, c : [p, 1] \to \mathbb{R}$ with $s_a(1-p) = s_b(2p-1) = s_c(1) = 0$ and $c(t) - a(t-p) = f(t) - f(t-p) - \delta$. It holds true for $t \in [p, 1]$ fixed

$$\begin{aligned}\min |a(t-p)|^2 + |c(t)|^2 &= |\frac{1}{2}(-c(t) + a(t-p))|^2 + |\frac{1}{2}(c(t) - a(t-p))|^2 \\ &= \frac{1}{2}|(c(t) - a(t-p))|^2 = \frac{1}{2}|f(t) - f(t-p) - \delta|^2,\end{aligned}$$

where the minimum is taken over all $a(t-p), c(t) \in \mathbb{R}$ with $c(t) - a(t-p) = f(t) - f(t-p) - \delta$. Hence,

$$\begin{aligned}\|\nabla_p s_f\|_{p-Sl}^2 =& \frac{1}{3p-1}(2s_f(p)^2 + 2(1-p)s_f(p)\delta + \delta^2(1-p)p) \\ &+ \min \|a(t)\|_{L^2[0,1]}^2 + \|c(t)\|_{L^2[0,1]}^2. \\ =& \frac{1}{3p-1}(2s_f(p)^2 + 2(1-p)s_f(p)\delta + \delta^2(1-p)p) \\ &+ \frac{1}{2}\|(f(t) - f(t-p) - \delta)_{t\in[p,1]}\|_{L^2[p,1]}^2 \\ =& \frac{1}{3p-1}(2s_f(p)^2 + 2(1-p)s_f(p)\delta + \delta^2(1-p)p) \\ &+ \frac{1}{2}(\|(f(t) - f(t-p))_{t\in[p,1]}\|_{L^2[p,1]}^2 - (1-p)\delta^2) \\ =& \frac{1}{2(3p-1)}(2s_f(p) + \delta(1-p))^2 \\ &+ \frac{1}{2}\|(f(t) - f(t-p))_{t\in[p,1]}\|_{L^2[p,1]}^2.\end{aligned}$$

This minimum is obtained at the function

$$\begin{aligned}f^*(t) =& \left(\frac{1}{3p-1}(s_f(p) + \frac{1-p}{2}\delta) + \frac{1}{2}(-f(t+p) + f(t))\right)\mathbf{1}_{[0,1-p]}(t) \\ &+ \frac{1}{2p-1}(s_f(p) - \frac{1-p}{3p-1}(s_f(p) - (2p-1)\delta))\mathbf{1}_{(1-p,p)}(t) \\ &+ \left(\frac{1}{3p-1}(s_f(p) + \frac{1-p}{2}\delta) + \frac{1}{2}(f(t+p) - f(t))\right)\mathbf{1}_{[p,1]}(t), t \in [0,1].\end{aligned}$$

Proof of Lemma 3.2 We have to prove Eq. (1) for the p-Slepian-process. To this end, let $p \le t \le 1, f \in L^2[0,1]$ and $(\nabla_p s_f)(\cdot) = s_f(p) + s_{\nabla_p f}(\cdot) \in \mathcal{H}_{p-\mathrm{Sl}}$. It

holds true for $t \in [p,1]$:

$$\begin{aligned}
&\mathbf{E}\left(\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]}(t)\cdot\frac{3s_f(p)+s_{\nabla_p f}(1)}{2(3p-1)}\left((\nabla_p B)_{[p,1]}(p)+(\nabla_p B)_{[p,1]}(1)\right)\right)\\
&\qquad+\mathbf{E}\left(\frac{1}{\sqrt{p}}(\nabla_p B)_{[p,1]}(t)\cdot\frac{1}{2}\int_p^1\nabla_p f(s)\,d(\nabla_p B)_{[p,1]}(s)\right)\\
&=\frac{3s_f(p)+s_{\nabla_p f}(1)}{2\sqrt{p}(3p-1)}\mathbf{E}\left((B_{[0,1]}(t)-B_{[0,1]}(t-p))(B_{[0,1]}(p)+B_{[0,1]}(1)-B_{[0,1]}(1-p))\right)\\
&\qquad+\frac{1}{2\sqrt{p}}\mathbf{E}\left((B_{[0,1]}(t)-B_{[0,1]}(t-p))\int_p^1\nabla_p f(s)\,d(B_{[0,1]}(s)-B_{[0,1]}(s-p))\right)\\
&=\frac{3s_f(p)+s_{\nabla_p f}(1)}{2\sqrt{p}}-\frac{1}{2\sqrt{p}}\left(s_{\nabla_p f}(1)-s_{\nabla_p f}(p)\right)=\frac{1}{\sqrt{p}}s_f(p)+s_{\frac{1}{\sqrt{p}}\nabla_p f}(t).
\end{aligned}$$

References

[1] Abrahams, J. (1984). Ramp crossings for Slepian's process. *IEEE Trans. Inform. Theory*, **30** No. 3, 574–575.

[2] Adler (1990). *An introduction to continuity, extrema, and related topics for general Gaussian processes*, Hayward, Calif., Institute of Mathematical Statistics, Lecture notes, monograph series 12.

[3] Bar-David (1975). Radon-Nikodym Derivatives, Passages and Maxima for a Gaussian Process with Particular Covariance and Mean. *Journal of Applied Probability*, **12**, 724–733.

[4] Bischoff, W. and Miller, F. (2000). Asymptotically optimal tests and optimal designs for testing the mean in regression models with applications to change–point problems. *Ann. Inst. Stat. Math.*, **52**, 658–679.

[5] Bischoff, W. and Hashorva, E. (2005). A lower bound for boundary crossing probabilities of Brownian bridge with trend. *Stat. Probab. Lett.*, **74**, 265–271.

[6] Cameron, R. H. and Martin, W. T. (1944). Transformations of Wiener Integrals Under Translations. *Annals of Mathematics*, **45**, 386–396.

[7] Chu, C., Hornik, K. and Kaun, C. (1995). MOSUM tests for parameter constancy. *Biometrika*, **82**, 603–617.

[8] Cressie, N. (1980). The Asymptotic Distribution of the Scan Statistic under Uniformity. *The Annals of Probability*, **8**, 828–840.

[9] Gao, F. and Li, W. (2007). Small ball probabilities for the Slepian Gaussian fields. *Trans. Am. Math. Soc.*, **359** No. 3, 1339–1350.

[10] Gegg, A. (2013). *Moving Windows zum Auffinden von Change-Points.* Doktor Thesis (in German), Catholic University of Eichstätt-Ingolstadt.

[11] Jamison (1970). Reciprocal processes: the stationary Gaussian case. *Ann. Math. Statist.* **41**, 1624-1630.

[12] Lifshits, M. (2012). *Lectures on Gaussian processes.* SpringerBriefs in Mathematics, Springer, Berlin.

[13] Liu, J. V., Zongfu, H. and Mao, H. (2014). Karhunen-Loève expansion for additive Slepian processes. *Stat. Probab. Lett.*, **90**, 93–99.

[14] Mörters, P. and Peres, Y. (2010). *Brownian motion.* Cambridge Series in Statistical and Probabilistic Mathematics 30, Cambridge University Press, Cambridge.

[15] Nikitin, Ya. Yu. and Orsingher, E. (2006). Sharp small ball asymptotics for Slepian and Watson processes in Hilbert norm. (Russian) *Zap. Nauchn. Semin. POMI*, **320**, 120–128 (2004), translation in *J. Math. Sci., New York*, **137** No. 1, 4555–4560.

[16] Shepp, L. A. (1966). Radon-Nikodym derivatives of Gaussian measures. *Ann. Math. Stat.*, **37**, 321–354.

[17] Shepp, L. A. (1971). First passage time for a particular Gaussian process. *Ann. Math. Stat.*, **42**, 946–951.

[18] Slepian, D. (1961). First passage time for a particular Gaussian process. *Ann. Math. Stat.*, **32**, 610–612.